

Transcriptional regulation by changes in tonicity

JOSEPH S. HANDLER and H. MOO KWON

Division of Nephrology, Johns Hopkins School of Medicine, Baltimore, Maryland, USA

Transcriptional regulation by changes in tonicity. Most organisms respond to a hypertonic environment by accumulating small organic solutes. In contrast to high concentrations of electrolytes, the small organic solutes do not perturb the activity of enzymes and other macromolecules within the cell. When the renal medulla becomes hypertonic during antidiuresis, multiple signaling pathways are activated. Here, we review the role of tonicity responsive enhancers (TonE) binding protein (TonEBP), a transcription factor activated in hypertonic cells. The activation of TonEBP by hypertonicity results from its translocation to the nucleus as well as an increase in TonEBP mRNA and protein. TonEBP may have a role beyond the response to tonicity since it is highly expressed in activated lymphocytes and in developing tissues.

The concentration of electrolytes within renal medullary cells remains remarkably constant despite large changes in the effective osmotic activity (tonicity) of the interstitium of the renal medulla (sodium varies between 140 mmol/L in hydrated rats and more than 400 mmol/L in antidiuretic rats) [1]. The stability of the intracellular electrolyte concentration results from the tonicity-regulated accumulation of specific small organic solutes that, in contrast to changing concentrations of electrolytes, do not perturb the activity of proteins and other macromolecules within cells [2]. The nonperturbing organic solutes that are accumulated by cells in the hypertonic medulla and in a variety of cultured cells are polyhydric alcohols (myoinositol and sorbitol), methylamines (betaine and glycerophosphorylcholine), and amino acids (taurine, alanine, proline, and to a lesser extent, other amino acids). Nonperturbing solutes are accumulated as the result of changes in the activity of rate-limiting, sodium-coupled transporters that carry the organic solutes into cells against steep concentration gradients (myoinositol, betaine, amino acids) and changes in activity of rate-limiting enzymes that catalyze solute synthesis (sorbitol) or degradation (glycerophosphorylcholine). Molecular cloning of the cDNAs for the rate-limiting proteins revealed that the increases in activities of those proteins

result from a hypertonicity-induced increase in the transcription of their genes and the resulting increase in synthesis of the proteins. The single exception is a transcription-independent decline in the activity of glycerophosphorylcholine phosphodiesterase, which degrades glycerophosphorylcholine [3]. Within the past few years, we have gained further insight into the molecular mechanism of the regulation of transcription by hypertonicity.

The regulation of transcription of most nonhousekeeping genes depends on specific DNA elements that are typically located in the region 5' to their promoter. Tonicity responsive enhancers (TonEs) have been identified in that region of the gene for the betaine transporter (BGT1) [4] and subsequently in that region of the gene for aldose reductase (AR), where they were designated osmotic response elements (ORE) [5–7], and in distant 5' regions of the SMIT gene [8]. The TonE consensus sequence TGGAAANNYYN confers transcriptional stimulation by hypertonicity on reporter genes with TonEs in their 5' region. In addition, the nucleus of hypertonic cells contains a protein that binds to radiolabeled TonEs in vitro in electrophoretic mobility shift assays (EMSAs) with far more activity than that found in the nucleus of isotonic cells. In hypertonic cells, specific G residues within the TonEs of BGT1 and SMIT are protected from methylation in so-called in vivo footprinting assays, whereas in isotonic cells, those G residues are methylated [8]. The changes in EMSA mobility and the protection from methylation in hypertonic cells suggest that the activation of transcription conferred by TonEs is the result of the binding of one or more proteins to TonEs.

Using TonEs as bait in yeast one-hybrid assays, we cloned the cDNA for a novel transcription factor, TonE binding protein (TonEBP), a large protein of approximately 160 kD [9]. Unexpectedly, mRNA for TonEBP is expressed in every tissue examined. The highest levels of expression are found in the renal medulla, brain, and heart. Figure 1 presents cartoon diagrams of TonEBP and the NFAT transcription factors, which play a key role in the regulation of transcription in activated lymphocytes, heart, and mesenchymal cells. The TonEBP sequence carboxy to the DNA binding domain contains many glutamine residues, a feature commonly found in

Key words: TonEBP, calcineurin, cyclosporine A, myoinositol, betaine, taurine, sorbitol.

© 2001 by the International Society of Nephrology




Fig. 1. Schematic comparison of the primary sequences of tonicity responsive enhancer binding protein (TonEBP) and the NFAT family. The grey boxes labeled “DB” denote DNA binding domains, while the striped boxes labeled “NHR” denote the NFAT homology region. NHR binds calcineurin and is dephosphorylated when calcineurin is activated. Note that NHR is not found in TonEBP. “QQ” and “QQQ” denote stretches of glutamine’s transcriptional regulation by changes in tonicity.

the activator region of transcription factors. The hypothesis that the glutamine-rich region mediates transcriptional activation was supported when we expressed a peptide lacking the presumed activator domain. The truncated protein behaved as a dominant negative; that is, it inhibited the transcriptional activation of reporter genes in hypertonic cells [9]. The DNA binding domain of TonEBP has a 45% identity to the DNA binding domains of the NFAT family of transcription factors, which share about 80% identity in that region. The greater identity within the DNA binding domain of NFATs includes a region that interacts with the transcription factor AP1. TonEBP lacks the region for interaction with AP1 and, in contrast to the NFAT transcription factors, stimulates transcription in the absence of activated AP1 [10]. Using a variety of strategies, other laboratories have subsequently cloned TonEBP and called it NFAT5 [10], NFATL1 [11], NFATz [12], and OREBP [13]. In contrast to the members of the NFAT family, TonEBP has no sequence homology to NFAT proteins outside of the DNA binding domain. Specifically, TonEBP lacks a sequence that is located amino terminal to the DNA binding domain, is phosphorylated by a nuclear protein kinase and directs the export of NFATs to the cytoplasm. In quiescent cells, NFATs are cytosolic. When lymphocytes are activated and cell calcium rises, NFATs are dephosphorylated by the calcium-calmodulin-sensitive phosphatase calcineurin and move to the nucleus where they regulate expression of specific genes. Activation of NFATs is inhibited by cyclosporine A and by FK506, which inhibit the phosphatase activity of calcineurin. Although there is a preliminary report of inhibition of the transcriptional response to hypertonicity by cyclosporine A and by FK506 using high concentrations of these agents (abstract; Sheikh-Hamad et al, *J Am Soc Nephrol* 11:47A, 2000), we found that cyclosporine A

and FK506 had no effect on the stimulation by hypertonicity of BGT1 activity and had divergent effects on the stimulation of SMIT activity [14]. Another laboratory found that concentrations of cyclosporine A or FK506 that clearly inhibited calcineurin activity in lymphocytes did not inhibit the transcriptional response to hypertonicity in the same cells [11].

TonEBP undergoes a number of changes when it is activated by hypertonicity. First, TonEBP is phosphorylated when cells are shifted to a hypertonic medium [15]. Most of the phosphorylation occurs at serine residues, but also there is increased tyrosine phosphorylation. Since TonEBP contains 197 serine residues and 13 tyrosine residues, it will be difficult to identify the important phospho-residues. Using immunofluorescence and immunoprecipitation from cell fractions, we found that in isotonic cells, approximately half of the cell’s TonEBP is in the cytoplasm and half in the nucleus. Within an hour of exposure to hypertonicity, cytoplasmic TonEBP diminishes, and nuclear TonEBP increases. After 8 to 12 hours, almost all of the TonEBP is nuclear [9]. In view of the similar time course for phosphorylation of TonEBP and its movement to the nucleus, we hypothesize that the phosphorylation plays a role in moving TonEBP to the nucleus, but this remains to be tested. In vitro binding assays showed that the association of TonEBP with TonE enhancer sequences is not affected by phosphorylation. Proteasome activity is also important in the regulation of TonEBP by hypertonicity. When proteasome protease activity is inhibited, movement of TonEBP to the nucleus is decreased, as is the increase in nuclear TonEBP activity [16]. In analogy to the regulation of nuclear factor- κ B (NF- κ B), there may be a cytosolic protein that limits TonEBP movement to the nucleus in control cells that is degraded by proteasomes in hypertonic cells, but for the moment this is only specula-

tion. In addition to translocation to the nucleus, activation by hypertonicity also evokes an increase in total cell TonEBP. The increase in TonEBP protein follows an increase in TonEBP mRNA abundance, which is probably the result of stimulating the transcription of TonEBP. Unfortunately, the very low level of transcription of TonEBP precludes assessment of transcription by classic nuclear run-off assays. The regulation of TonEBP by tonicity is bidirectional. When cells are switched from a hypertonic to an isotonic medium, the degradation of TonEBP mRNA and protein is not affected by tonicity, as they decay with half-lives of approximately 6 and 10 hours, respectively. Since TonEBP has a detectable level of activity in isotonic cells, we examined its status in cells adapted to growth in medium made hypotonic by removing most of the NaCl in the medium. The level of TonEBP fell in hypotonic cells, and nuclear immunostaining decreased far more than in cytosolic cells [17]. We return to these observations later when speculating about the regulation and function of TonEBP.

There is a large gap in understanding how hypertonicity regulates TonEBP. First, how is hypertonicity detected? The strategy of maintaining intracellular ionic activity nearly constant by accumulating or losing organic osmolytes as external tonicity changes is an old one. Like the animal cells we study, bacteria, yeast, and most plant and animal cells accumulate amino acids and polyhydric alcohols when exposed to a hypertonic environment [2]. Some bacterial osmolyte transporters are directly stimulated by hypertonicity [18, 19]. The initial sensor in yeast is a two-component histidine kinase located in the plasma membrane [20]. The sensor initiates a multistep phosphorelay cascade that activates a p38 mitogen-activated protein (MAP) kinase homologue, HOG kinase, which regulates the rate of transcription of a number of genes. Similar sensors have not been identified in animal cells, although there is evidence that animal cells detect changes in cell size or shape within minutes after exposure to a hypertonic environment [21, 22]. Since most cells return to near their original isotonic size within 30 minutes as a result of a regulatory volume increase mediated by ion uptake, it is unlikely that the activation of TonEBP, which occurs over hours, is mediated by cell size or shape. Multiple changes, including the rapid induction and suppression of many genes, occur when cells are abruptly shifted to a hypertonic environment. The cell cycle is arrested, and many regulatory pathways are activated or inhibited [23, 24]. Indirect evidence suggests that the sensor that regulates activation of TonEBP may detect intracellular ion activity. The activation of transcription by hypertonicity is reduced as cells accumulate organic osmolytes and cell ion concentrations return to those of isotonic cells. In cells in hypertonic medium, there is a close correlation between the increase in total cell cation concentration and the increase in activity of

AR [25]. Also, inhibition or stimulation of osmolyte accumulation by the removal or addition of transporter substrate prolongs or hastens the decline in the abundance of hypertonicity-induced taurine transporter [26] or AR mRNA [27].

There are many unanswered questions regarding TonEBP. TonEBP mRNA is expressed at low levels in most cells in culture and in virtually all tissues that have been sampled. However, only the kidney medulla normally experiences marked changes in tonicity. Does TonEBP regulate processes other than those involved in the response to hypertonicity? The low level of expression of TonEBP in isotonic cells is reduced further when cells are adapted to growth in hypotonic medium, as though a hypotonic state is the null condition. Perhaps the levels of TonEBP expression in isotonic cells are based on life in more hypotonic environments experienced by ancestral precursor cells. A recent report that TonEBP, like some NFAT transcription factors, is induced in lymphocytes by ConA or by ionomycin plus a phorbol ester is intriguing [11]. In addition, TonEBP is expressed at much higher levels in embryonic tissues than in adult tissues [11]. Does TonEBP play a role in development?

ACKNOWLEDGMENT

The work from this laboratory was supported by National Institutes of Health Grants DK4448s and DK42479.

Reprint requests to Joseph S. Handler, M.D., 965 Ross Research Building, Johns Hopkins School of Medicine, 720 Rutland Avenue, Baltimore, Maryland 21205, USA.
E-mail: jhandler@jhmi.edu

REFERENCES

1. BECK F, DORGE A, RICK R, THURAU K: Intra- and extracellular element concentrations of rat renal papilla in antidiuresis. *Kidney Int* 25:397–403, 1984
2. YANCEY PH, CLARK ME, HAND SC, *et al*: Living with water stress: Evolution of osmolyte system. *Science* 217:1214–1222, 1982
3. KWON ED, ZABLOCKI K, JUNG KY, *et al*: Osmoregulation of GPC:choline phosphodiesterase in MDCK cells: Different effects of urea and NaCl. *Am J Physiol* 269:C35–C41, 1995
4. TAKENAKA M, PRESTON AS, KWON HM, HANDLER J: The tonicity-sensitive element that mediates increased transcription of the betaine transporter gene in response to hypertonic stress. *J Biol Chem* 269:29379–29381, 1994
5. FERRARIS JD, WILLIAMS CK, JUNG K-Y, *et al*: ORE, a eukaryotic minimal essential osmotic response element: The aldose reductase gene in hyperosmotic stress. *J Biol Chem* 271:18318–18321, 1996
6. DAOUDAL S, TOURNAIRE C, HALERE A, *et al*: Isolation of the mouse aldose reductase promoter and identification of a tonicity-responsive element. *J Biol Chem* 272:2615–2619, 1997
7. KO BCB, CHUNG SM: Identification and characterization of multiple osmotic response sequences in the human aldose reductase gene. *J Biol Chem* 272:16431–16437, 1997
8. RIM JS, ATTA MG, DAHL SC, *et al*: Transcription of the sodium/myo-inositol cotransporter gene is regulated by multiple tonicity-responsive enhancers spread over 50 kilobase pairs in the 5'-flanking region. *J Biol Chem* 273:20615–20621, 1998
9. MIYAKAWA H, WOO SK, DAHL SC, *et al*: Tonicity-responsive enhancer binding protein, a Rel-like protein that stimulates transcrip-

- tion in response to hypertonicity. *Proc Natl Acad Sci USA* 96:2538–2542, 1999
10. LOPEZ-RODRIGUEZ C, ARAMBURU J, RAKEMAN AS, RAO A: NFAT5, a constitutively nuclear FNAT protein that does not cooperate with Fos and Jun. *Proc Natl Acad Sci USA* 96:7214–7219, 1999
 11. TRAMA J, LU Q, HAWLEY RG, HO SN: The NFAT-related protein NFATL1 (TonEBP/NFAT5) is induced upon T cell activation in a calcineurin-dependent manner. *J Immunol* 165:4884–4894, 2000
 12. PAN S, TSURITA R, MASUDA ES, *et al*: NFATz: A novel rel similarity domain containing protein. *Biochem Biophys Res Commun* 272:765–776, 2000
 13. KO BCB, TURCK CW, LEE KW, *et al*: Purification, identification, and characterization of an osmotic response element binding protein. *Biochem Biophys Res Commun* 270:52–61, 2000
 14. ATTA MG, DAHL SC, KWON HM, HANDLER JS: Tyrosine kinase inhibitors and immunosuppressants perturb the myo-inositol but not the betaine cotransporter in isotonic and hypertonic MDCK cells. *Kidney Int* 55:956–962, 1999
 15. DAHL SC, HANDLER JS, KWON HM: Hypertonicity-induced phosphorylation and nuclear localization of the transcription factor TonEBP. *Am J Physiol (Cell Physiol)* 280:C248–C253, 2001
 16. WOO SK, MAOYUO D, HANDLER J, KWON HM: Nuclear redistribution of tonicity-responsive enhancer binding protein requires proteasome activity. *Am J Physiol (Cell Physiol)* 278:C323–C330, 2000
 17. WOO SK, DAHL SC, HANDLER JS, KWON HM: Bidirectional regulation of tonicity-responsive enhancer binding protein in response to changes in tonicity. *Am J Physiol (Renal Physiol)* 278:F1006–F1012, 2000
 18. RACHER KI, VOEGELE RT, MARSHALL EV, *et al*: Purification and reconstitution of an osmosensor: Transporter ProP of *Escherichia coli* senses and responds to osmotic shifts. *Biochem J* 38:1676–1684, 1999
 19. HUANG K-J, LAN C-Y, IGO MM: Phosphorylation stimulates the cooperative DNA-binding properties of the transcription factor OmpR. *Proc Natl Acad Sci USA* 94:2828–2832, 1997
 20. POSAS F, WURGLER-MURPHY SM, MAEDA T, *et al*: Yeast HOG1 MAP kinase cascade is regulated by a multistep phosphorelay mechanism in the SLN1–YPD1–SSK1 “two-component” osmosensor. *Cell* 86:865–875, 1996
 21. SZASZ K, BUDAY L, KAPUS A: Shrinkage-induced protein tyrosine phosphorylation in Chinese hamster ovary cells. *J Biol Chem* 272:16670–16678, 1997
 22. KRUMP E, NIKITAS K, GRINSTEIN S: Induction of tyrosine phosphorylation and Na⁺/H⁺ exchanger activation during shrinkage of human neutrophils. *J Biol Chem* 272:17303–17311, 1997
 23. DMITRIEVA N, KULTZ D, MICHEA L, *et al*: Protection of renal inner medullary epithelial cells from apoptosis by hypertonic stress-induced p53 activation. *J Biol Chem* 275:18243–18247, 2000
 24. KULTZ D, MADHANY S, BURG MB: Hyperosmolality causes growth arrest of murine kidney cells induction of GADD45 and GADD153 by osmosensing via stress-activated protein kinase 2. *J Biol Chem* 273:13645–13651, 1998
 25. UCHIDA S, GARCIA-PEREZ A, MURPHY H, BURG MB: Signal for induction of aldose reductase in renal medullary cells by high external NaCl. *Am J Physiol (Cell Physiol)* 256:C614–C620, 1989
 26. UCHIDA S, KWON HM, YAMAUCHI A, *et al*: Molecular cloning of the cDNA for an MDCK cell Na⁺- and Cl⁻-dependent taurine transporter that is regulated by hypertonicity. *Proc Natl Acad Sci USA* 89:8230–8234, 1992
 27. MORIYAMA T, GARCIA-PEREZ A, OLSON AD, BURG MB: Intracellular betaine substitutes for sorbitol in protecting renal medullary cells from hypertonicity. *Am J Physiol (Renal Fluid Electrol)* 260:F494–F497, 1991